arXiv:cond-mat/9909074v1 [cond-mat.str-el] 6 Sep 1999

The Ground State of the Two-Dimensional Hubbard Model

Yoshihiro Asai
Physical Science Division, Electrotechnical Laboratory (ETL),
Agency of Industrial Science and Technology (AIST),
Umezono 1-1-4, Tsukuba, Ibaraki 305-8568, Japan

(Received on February 1, 2008)

We have studied the ground state of the two-dimensional Hubbard model by using the recently proposed adaptive sampling quantum monte carlo (ASQMC) method. We have paid attention to the model whose non-interacting band dispersion is almost flat near $(\pi, 0)$. To minimize the effect of the finite size gap overlying on the Fermi level, we have tuned both filling and band structure. We found enhancement of the d-wave correlation function at large distance, the spin gap and the momentum distribution function consistent with the d-wave gap. We also found the coexistence of both the commensurate and incommensurate peaks in $S(\vec{q})$, which does not contradict a recent experimental finding that both the resonance peak and the incommensurate peaks reside in the same doping level of YBCO and BSCCO.

It is very difficult to calculate ground state properties of large two-dimensional (2D) Hubbard clusters away from half filling when U/t is not small because of the notorious negative sign problem for quantum monte carlo (QMC) methods. We have recently proposed the adaptive sampling quantum monte carlo (ASQMC) method to cope with the problem. [1] The start of the project to develop the ASQMC method was influenced by the publication of the constrained path (CP) monte carlo method by Zhang, Carlson and Gubernatis (ZCG). [2] However we found it is possible to reduce the negative sign ratio without any constraint if we adopt a new update scheme. This may be possible because the update scheme adopted by ZCG itself seems to suppress the negative sign ratio, which should be the reason why their method is superior in accuracy than the more elaborate positive projection method proposed by Fahy and Hammann 8 years ago. [3] We developed our update scheme within the frame work of the "exact update method" of the auxiliary field quantum monte carlo method. Our scheme is not only easier to understand for those who have been working on Hubbard and related models but also it is superior in accuracy than the ZCG's scheme because it is free from the inaccurate mixed estimator method for measurements and its variants which are indispensable to the CP method. This should be one of the main reasons why our ASQMC method gives more accurate results than the CP method. Actually, we observed the ASQMC gives better value for the ground state energy of 14 electron 4×4 cluster when $U/t = 12$. [1] Even without constraint, the negative sign ratio is reduced largely in the all cases studied, which makes measurements stable. $< Sign >$ decrease to 0 very slowly as we increase the projecting imaginary time τ. (Fixing $\Delta\tau$ and τ_c constant.) Typically, $\tau = 20$ to 30 is feasible. We have made the serial correlation test in many cases and found that samples obtained by the ASQMC are statistically independent and the variance calculated by the standard formula gives a good estimate for the true variance. As an example we show the autocorrelation function for the d-wave superconducting correlation function at $\vec{q} = 0$ in the case of parameter set (i) described below in Fig. 1.

We used the ASQMC method to study the ground state properties of the 2D Hubbard model. The parameter region we paid our attention was such that the non-interacting band dispersion was almost flat near the $(\pi, 0)$ point. We tuned electron filling and band dispersion such that the energy difference between the highest occupied level (HOL) and the lowest unoccupied level (LUL) was close to zero. The Hubbard model we studied may belong to different class than that frequently studied on the bipartite lattice. We have studied various parameter regions of 6×6, 8×8, 10×10 and 12×12 clusters. [4] Here we discuss following three cases; (i) 6×6 lattice with $t' = -0.1667$, and $t'' = 0.2$. $U = 4$. The number of electron is 28. (ii) 10×10 lattice with $t' = -0.2$, and $t'' = 0.05$. $U = 4$. The number of electron is 84. (iii) 12×12 lattice with $t' = -0.1676$, and $t'' = 0.2$. $U = 4$. The number of electron is 92. To make sure that our result is convergent to the $\tau \to \infty$ limit, we show the τ dependence of the ground state energy of (i) in Fig. 2. The calculation was stable at least up to $\tau = 20$, but the energy converged at $\tau = 3$. We observed similar fast convergence in the case of (ii). Hence, we proceeded calculations with $\tau = 3$ to save CPU time. The fast convergence suggests that energy gaps among the states with the same symmetry are relatively large, while the gap between the ground and low lying excited states in the finite size system can be small, if these state have different symmetries. This is because we are using the projector method. We found the enhancement of the d-wave superconducting correlation at large distance in the ground state when we used the parameter sets (i) and (ii). We also found the momentum distribution function can be well described by using the d-wave BCS mean field theory. We found exponential decay of the distance dependence of the spin correlation function, which is suggestive of opening the spin gap. [4] All these are consistent with the d-wave superconducting ground state. Even more surprisingly, the commensurate peak coexists with the incommensurate peaks in $S(\vec{q}) = \int d\omega S(\vec{q}, \omega)$ of the superconducting state. This is characteristic to the parameter region involving the three parameter sets. The coexistence was most clearly seen in a over doped region (iii) which was shown in Fig. 3. (The peak interval between the incommensurate peaks is largest in (iii).) Experiments on YBCO and BSCCO seem to suggest that both the resonance peak and the incommensurate peaks exist at different ω but in the same filling. [5] *The suggestion from experiments does not contradict our result on* $S(\vec{q})$. In the other parameter regions, we did not so far observe similar $S(\vec{q})$. The numerical results so far obtained by using the ASQMC method suggest that the band dispersion close to $(\pi, 0)$ seems to be very crucial to the ground state of the 2D Hubbard model.

---

[1] Y. Asai, cond-mat/9904255.
[2] S. Zhang, J. Carlson, and J.E. Gubernatis, Phys.Rev.B55, (1997) 7464.
[3] S.B. Fahy and D.R. Hamman, Phys.Rev.B43, (1991) 765.
[4] Y. Asai, cond-mat/9904310.
[5] H.A. Mook, F. Doğan, and B.C. Chakoumakos, cond-mat/9811100.



FIG. 1. Autocorrelation function for the d-wave superconducting correlation at $\vec{q} = 0$ in the case of the parameter set (i).



FIG. 2. The projecting imaginary time τ dependence of the ground state energy of the 6×6 cluster.



FIG. 3. The spin correlation function $S(\vec{q})$ of the 12×12 cluster.